

**PROPOSAL FOR INCLUSION OF SPECIES ON THE APPENDICES OF THE
CONVENTION ON THE CONSERVATION OF MIGRATORY SPECIES OF
WILD ANIMALS**

- A. PROPOSAL:** Change listing in Appendix II of *Tursiops truncatus* from “western Mediterranean population” to “Mediterranean population”
- B. PROPONENT:** Principality of Monaco
- C. SUPPORTING STATEMENT:**

1. Taxon

1.1 Classis	Mammalia
1.2 Ordo	Cetacea
1.3 Familia	Delphinidae
1.4 Species	<i>Tursiops truncatus</i> (Montagu, 1821)
1.5 Common name(s)	English: Bottlenose dolphin French: Grand dauphin, Dauphin souffleur Spanish: Delfín mular Albanian: Delfin i madh Arabic: دلفين كبير (Delfin kabir) Croatian: Dobri dupin (Dolphinan yam hatichon) Hebrew: דולפין - דולפין Italian: Tursiope Maltese: Delfin geddumu qasir Turkish: Afalina

2. Biological data

2.1 Distribution

Bottlenose Dolphins have been reported to mostly occur the coastal waters of the all Mediterranean basin. They are considered regular off Algeria (Boutiba *et al.*, 2003), Croatia (Bearzi *et al.*, 1997), Cyprus (Hadjichristoforou, 2004), France (Deguy & Cyrus 1973; Duguay *et al.*, 1983; Ripoll *et al.*, 2004; Dhermain, 2006), Greece (Frantzis *et al.*, 2003), Israel (Feingold *et al.*, 2005; Scheinin *et al.*, 2005), Italy (Notarbartolo *et al.*, 1993), Morocco (Bayed, 1997), Slovenia (Genov & Kotnjek, 2007), Spain (Cañadas *et al.*, 2002; Raga & Pantoja, 2004; Cañadas & Hammond, 2006), Tunisia (Chakroun, 1994; Ben Naceur *et al.*, 2004) and Turkey (Öztürk *et al.*, 2004).

The Bottlenose Dolphin is one of the most common cetacean species in the Mediterranean, after the Striped Dolphin *Stenella coeruleoalba*, particularly in continental shelf waters (Bearzi *et al.*, 2008). Bottlenose Dolphins are highly adaptable and inhabit a wide variety of habitats including lagoons and enclosed seas (Bearzi & Ferretti 2000; Bearzi *et al.*, 2007a), deep areas with steep bottom gradients (e.g. around Crete; Frantzis *et al.*, 2003), productive waters 200-500 m deep (e.g. in the Alborán Sea; Cañadas & Hammond, 2006), and the

channels and shelf waters of archipelagos (Pulcini *et al.*, 1993; Bearzi *et al.*, 1997; Impetuoso *et al.*, 2003; Mussi & Miragliuolo, 2003; Forcada *et al.*, 2004). Many of the Mediterranean areas inhabited by these dolphins are subject to intensive human use, e.g. the straits of Gibraltar, Bonifacio, and Messina (Romeo *et al.*, 2003; Raga & Pantoja 2004; Dhermain, 2006), and the gulfs of Lion, Genova, and Trieste (Francese *et al.*, 1999; Ripoll *et al.*, 2004; Gnone *et al.*, 2006). Bottlenose Dolphins are also found around oil and gas drilling platforms in the Adriatic Sea (Triossi & Tizzi, 2003). Within their overall range, gaps with very low densities of animals have been documented, e.g. in the north-western Ligurian Sea (France and Italy) and in the north-western Gulf of Vera (Spain). Variation in density is likely related to several factors, including a) habitat characteristics, b) local availability of suitable prey, and c) the generally gregarious nature of Bottlenose Dolphin communities. Moreover, the effects of past extermination campaigns (Bearzi *et al.*, 2004a) and a variety of ongoing threats probably have contributed to the pattern of present-day occurrence of Bottlenose Dolphins across the region.

See section 2.3 (Habitat) for more details.

2.2 Population

2.2.1 Numbers: absolute and relative abundances

Little is known about the numbers of Bottlenose Dolphins in the Mediterranean Sea. There is no basin-wide estimate. The most reliable information comes from local studies conducted in the Strait of Gibraltar (Pérez *et al.*, 2006), Alborán Sea (Cañadas, 2006; Cañadas & Hammond, 2006), the Spanish waters between Gulf of Valencia and Gulf of Vera (Gómez de Segura *et al.*, 2006), Balearic Sea (Forcada *et al.*, 2004), a small portion of the Italian waters (Lauriano *et al.*, 2003), the Tunisian plateau (Ben Naceur *et al.*, 2004), the northern Adriatic Sea (Fortuna *et al.*, 2000), Maltese waters (Vella, 1999), portions of the Turkish and Greek seas (Bearzi *et al.*, 2007a). These numbers are based on different methodological approaches, including absolute abundance estimates obtained through mark-recapture methods or Generalized Additive Models (GAMs), maximum number of photo-identified individuals, and discovery curves (i.e. curves showing the cumulative number of individual dolphins identified as a function of photo-identification effort, usually expressed as survey days with photos). Additional information that could in future increase the number of available absolute estimates for different areas exist for Lampedusa Island (Italy; Pulcini *et al.*, 2004), northern Adriatic Sea (Bearzi *et al.*, 2007b; Genov & Fortuna, 2005), Central Adriatic Sea (Kornati and Murtar Sea, Croatia; Impetuoso *et al.*, 2003), Eastern Ionian Sea (Greece; Bearzi *et al.*, 2005, 2006), Amvrakikos Gulf (Greece; Bearzi *et al.*, 2007a), Ligurian Sea and northern Tyrrhenian Sea (Italy; Gnone *et al.*, 2006), Gulf of Catania (Italy; Tringali *et al.*, 2004), Waters off Corsica (France; Dhermain 2006), Eastern Ionian Sea (Greece; Bearzi *et al.*, 2005), Mediterranean waters of Israel (Scheinin *et al.*, 2005). Most studies in coastal waters are limited to relatively small areas of 400 – 1,000 km² and likely do not cover the entire range of the groups under study. Relatively recent, broad-scale shipboard surveys (4,000 – 80,000 km²) showed that in some Mediterranean areas Bottlenose Dolphins are present both near shore and offshore, and densities can range between 4 and 20 animals per 100 km² (Ben Naceur *et al.*, 2004; Forcada *et al.*, 2004; Cañadas & Hammond, 2006; Gómez de Segura *et al.*, 2006). In this situation, the total population size in the Mediterranean remains uncertain, but it is unlikely to exceed the low 10 000s (Bearzi & Fortuna, 2006).

Little information exists for other parts of the Mediterranean basin where only information on past and present occurrence comes from stranding records. Virtually nothing is known on abundances for large portions of the south-eastern part of the basin.

2.2.2 Population structure

Based on nuclear and mitochondrial DNA analyses, Mediterranean Bottlenose Dolphins resulted as genetically differentiated from those inhabiting the contiguous eastern North Atlantic Ocean and the Black Sea (Natoli *et al.*, 2005). The genetic analysis of 74 samples collected along a continuous distributional range from the Mediterranean Sea showed some degree of population structure with boundaries that coincide with the transitions between habitat regions. These regions can be characterized by ocean floor topography and by features such as surface salinity, productivity and temperature (Natoli *et al.*, 2005).

2.3 Habitat

Bottlenose Dolphins in the Mediterranean are often regarded as predominantly ‘coastal’ or ‘inshore’ animals but this designation may be misleading as they can be encountered in continental shelf and shallow plateau waters at any distance from the coast of either main land and islands (Notarbartolo di Sciara *et al.*, 1993; Bearzi *et al.*, 1997; Triossi & Tizzi 2003; Bearzi *et al.*, 2004a; Ben Naceur *et al.*, 2004; Cañadas *et al.*, 2004; Gómez de Segura *et al.*, 2004; Ripoll *et al.*, 2004; Gannier, 2005), but also in straits, gulfs, eutrophic waters of estuaries and lagoons steep coasts with no continental shelf and deep waters of the continental slope (Cañadas *et al.*, 2002; Bearzi & Ferretti 2000; Frantzis *et al.*, 2003; Zafiroopoulos & Merlini, 2003; Cañadas *et al.*, 2004; Forcada *et al.*, 2004; Gómez de Segura *et al.*, 2004; Bearzi *et al.*, 2005, 2007a). This species usually favours depths shallower than 200 m (Pace *et al.*, 1999; Mussi *et al.*, 1998; Gazo *et al.*, 2004b; Gnone *et al.*, 2006; Gonzalvo *et al.*, 2004; Manoukian *et al.*, 2004).

2.4 Migrations

Even though bottlenose dolphins can display rather strong site fidelity, they showed to have quite wide ranges in European waters, up to 400 km (Wilson *et al.*, 2004). In the Mediterranean Sea, coast-to-coast movements across open seas occur covering over 200 km, in Ligurian Sea, between Corse and Italy (Dhermain *et al.*, 199), in the Adriatic Sea, between Slovenia, Croatia and Italy (Fortuna 2006). In addition, transnational movement are very likely to occur between the continental waters of Lampedusa Island (Italy) and Tunisia (Pulcini *et al.*, 2004; Ben Naucer *et al.*, 2004), the eastern Aegean waters between Greece and Turkey (Frantzis *et al.*, 2003).

3. **Threat data**

Owing to their occurrence in coastal waters, bottlenose dolphins in the Mediterranean are exposed to a wide variety of human activities. Whilst intentional killing was likely the most important cause of mortality until the 1960s (see previous section), important ongoing threats include incidental mortality in fishing gear and the reduced availability of key prey caused by region-wide overfishing and environmental degradation. Additional potential or likely threats include the toxic effects of xenobiotic chemicals, epizootic outbreaks, direct disturbance from boating and shipping, noise, and the consequences of climate change. It is worth noting that this same array of known and potential threats applies to riverine, estuarine and coastal

cetaceans (and other groups of organisms) in many other parts of the world as well (e.g. Reeves *et al.*, 2003).

3.1 Direct threat of threat of the population

3.1.1 Illegal trade

The level of illegal trade is unknown. At present, live capture of bottlenose dolphins is prohibited in all Black Sea countries except for Turkey where permits for the live capture of 30 animals in the Black, Marmara, Aegean and Mediterranean Seas were issued and realized at least in part (23 captures were reported) during 2006 and 2007 (Convention on the Conservation of European Wildlife and Natural Habitats 2007).

3.1.2 Direct catch, captive breeding for commercial purposes (outside the country of origin) Direct catches were common in the past throughout the entire Mediterranean Sea, mainly in the framework of national fishery management schemes (Gourret, 1894; Barone, 1895; Del Rosso, 1905; Peksider-Srica, 1931; Brunelli, 1932; Crnkovic, 1958; Cuculic, 1960; Marelic, 1961; Duguay *et al.*, 1983; Bompar, 2000; Bearzi *et al.*, 2004a). Dolphins were considered detrimental for fisheries in terms of competition for the same resources, but in some area they were also considered as target species (Bearzi *et al.*, *in press*). The overall frequency of intentional killing has been drastically declining over the long term due to new legislation granting cetacean protection in most Mediterranean countries. At the present there are not known direct catches; however killing in retaliation for damage to fisheries, killing with harpoons or guns for local consumption of meat have been reported in the past in Ligurian and Tyrrhenian seas, notwithstanding legal protection (Di Natale, 1990; Di Natale & Notarbartolo di Sciara, 1994). Such events seem to have become extremely rare in recent times, but they may still occur.

3.1.3 Incidental catch and commercial exploitation

Due to their opportunistic behaviour and predominantly coastal occurrence, Bottlenose Dolphins in the Mediterranean are often exposed to entanglement in many types of fishing gear, in most of the Mediterranean countries, including Algeria, Croatia, France, Greece, Israel, Italy, Malta, Morocco, Spain, Tunisia, and Turkey (Di Natale, 1990, 1995; Consiglio *et al.*, 1992; Silvani *et al.*, 1992; Di Natale & Notarbartolo di Sciara, 1994; Öztürk *et al.*, 2001; Boutiba *et al.*, 2003; Raitsos *et al.*, 2003; Roditi-Elasar *et al.*, 2003; Tudela *et al.*, 2005; Kent *et al.*, 2005; Brotons *et al.*, 2006; Díaz López, 2006b; Fortuna, 2006; Van Canneyt & Peltier, 2006). Large pelagic driftnets (Di Natale & Notarbartolo di Sciara, 1994; Öztürk *et al.*, 2001), small pelagic driftnets (Di Natale & Notarbartolo di Sciara, 1994), set nets (Silvani *et al.*, 1992; Di Natale & Notarbartolo di Sciara, 1994; Raitsos *et al.*, 2003; Brotons *et al.*, 2006; Díaz López, 2006b), bottom trawls (Silvani *et al.*, 1992; Goffman *et al.*, 2001; Feingold *et al.*, 2005; Brotons *et al.*, 2006), purse seines (Bradai, 2001), traditional tuna traps (Di Natale & Notarbartolo di Sciara, 1994) and antipredator nets used in aquaculture facilities (Díaz Lopez & Bernal Shirai, 2007). In addition to incidental mortality, depredation and damage caused by dolphins to fishing gear may result in animals being shot or harassed in retaliation (Di Natale, 1990; Silvani *et al.*, 1992; Di Natale & Notarbartolo di Sciara, 1994; Commission of the European Communities, 2002; Fernández-Contreras *et al.*, 2002; Gazo *et al.*, 2004a; Mitra *et al.*, 2004; Blasi & Pace, 2006).

Few attempts have been made to estimate fishery-related mortality for Bottlenose Dolphins (Silvani *et al.*, 1992; Di Natale, 1995; Díaz Lopez, 2006b; Díaz Lopez & Bernal Shirai, 2007) and the magnitude of bycatch and retaliation events is unknown in most cases. Even when

they are available, bycatch estimates are partial in terms of geographic and gear coverage. However, the available studies and circumstantial evidence for local populations raise serious concern, suggesting that annual fishery-induced mortality is locally unsustainable in at least some cases (Silvani *et al.*, 1992; Brotons *et al.*, 2006; Díaz Lopez, 2006b; Fortuna 2006; Díaz Lopez & Bernal Shirai, 2007). Rigorous studies of bycatch rates using reliable methods, which normally must include on-board observers and a statistically robust sampling design, are needed to obtain credible estimates of mortality. It is then incumbent to determine 'sustainability' by reference to the population size, also taking into account other existing threats.

In recent years, takes due to illegal fisheries has been reported off Morocco (Tudela, 2004), Spain (Silvani *et al.*, 1992; Gazo *et al.*, 2004a; Tudela, 2004), Italy (Consiglio *et al.*, 1992; Blasi & Pace, 2006) and Greece (Mitra *et al.*, 2004).

The persisting illegal use of dynamite for fishing in several Mediterranean areas, including Algeria, Croatia, France, Greece, Lebanon, Libya and Malta (Di Natale, 1990; Reynolds *et al.*, 1994; Tudela, 2004; Fortuna, 2006; Notarbartolo di Sciara *et al.*, 2006; Dhermain & Cesarini, 2007), represents another fishery-related threat to Bottlenose Dolphins. Though impact at the basin level is probably low, it may be significant locally (Fortuna 2006).

3.2 Habitat destruction

3.2.1 Lack of food resources

Overlap between dolphin prey species and fishery target species does not necessarily imply direct competition (Briand, 2004). However, it is reasonable to infer competitive interactions of some kind when key prey become scarce and remain subject to heavy fishing pressure (Trites *et al.*, 1997). In this regard, we note that about 95% of marine-fish catches globally come from continental shelf regions (Roberts & Hawkins, 1999) where Bottlenose Dolphins occur. Overfishing is having profound direct and indirect impacts on Mediterranean ecosystems (Sala, 2004). In the Mediterranean there is an acute lack of historical data and fisheries statistics are generally incomplete and unreliable, data on fishing effort being almost absent (Briand, 2000, 2003; Leonart, 2005). Nonetheless, it is generally acknowledged that unsustainable fishing has contributed significantly to dramatic ecological changes and caused the decline of many fish stocks (Caddy & Griffiths, 1990; De Walle *et al.*, 1993; Stanners & Bourdeau, 1995; Caddy, 1997). According to FAO, approximately 35% of the Mediterranean stocks are exploited beyond MSY levels, and 43% at MSY levels (Garcia *et al.*, 2005). Some of the Mediterranean fish stocks that have been either 'overexploited' or 'fully exploited' include important Bottlenose Dolphin prey such as European Hake, Striped Red Mullet, European Pilchard, Common Pandora *Pagellus erythrinus*, Annular Seabream *Diplodus annularis*, and Atlantic Horse Mackerel *Trachurus trachurus* (Leonart, 2005).

Yet, the fact that availability of Bottlenose Dolphin prey is strongly influenced by fishing pressure, with potential consequences on dolphin density, status and population trends is rarely taken into due consideration. Reduced carrying capacity due to overfishing was proposed as one explanation for the low densities of Bottlenose Dolphins in the Adriatic and Ionian Seas (Bearzi *et al.*, 1999, 2005a, 2006). Conversely, dolphin densities tend to be high in areas where prey is still abundant. For instance, dolphin density in the prey-rich Amvrakikos Gulf, Greece - where effective fishery management measures including the prohibition of purse seining and trawling are in place - is one order of magnitude higher than in the overfished waters of the nearby island of Kalamos (Bearzi *et al.*, 2006, 2007a; Table 2).

3.2.2 Disease

Epizootic outbreaks appear to have affected Bottlenose Dolphins to a lesser extent than other Mediterranean delphinids, such as the Striped Dolphin (Aguilar & Raga, 1993; Van Bresse *et al.*, 1993). Morbillivirus infections have been reported in one individual Bottlenose Dolphin stranded on the Mediterranean coast of Israel in 1994 (Tsur *et al.*, 1997), and one stranded in Mauritania (Atlantic coast of West Africa) in 1988 (Van de Bildt *et al.*, 2001). However, Bottlenose Dolphins elsewhere have experienced mass mortality from such outbreaks, e.g. in Black Sea waters (Birkun *et al.*, 1998) and on the Atlantic coast of the United States, where more than half of one local population may have died (Lipscomb *et al.*, 1994; Duignan *et al.*, 1996; Schulman *et al.*, 1997). As epizootic phenomena may be related to immune-system compromise induced by exposure to xenobiotics and/or by stress from poor nutrition (Aguilar & Borrell, 1994; Calzada *et al.*, 1996; O'Shea & Aguilar, 2001), the risk of disease outbreaks in Bottlenose Dolphins in the Mediterranean may be considerable.

Toxoplasmosis that can cause mortality in many species of domestic and wild animals (Dubey *et al.*, 2003), were also found in specimen of bottlenose dolphins in Mediterranean Sea (Di Guardo *et al.*, 1995a, 1995b; Cabezon *et al.*, 2004).

3.3 Indirect threat

Toxic contamination is a major concern in marine mammal populations because of the potential effects on reproduction health and DNA damage (Gauthier *et al.*, 1999; O'Shea *et al.*, 1999; Fossi & Marsili, 2003; Newman & Smith 2006). The Bottlenose Dolphins' worldwide distribution and great adaptability to diverse habitats make this species a good indicator of the quality of inshore marine ecosystems.

Contaminant levels, particularly of organochlorine compounds, in Mediterranean Bottlenose Dolphins are very high compared to the levels reported for Bottlenose Dolphins in other areas (Corsolini *et al.*, 1995; Marsili & Focardi, 1997; Aguilar *et al.*, 2002; Fossi & Marsili, 2003; Wafo *et al.*, 2005; Borrell *et al.*, 2006; Borrell & Aguilar, 2007). At concentrations similar to or lower than those documented for Mediterranean Bottlenose Dolphins, compounds such as PCBs or PAHs have been associated with reproductive disorders, immune-system suppression, and neoplasia in other populations of Bottlenose Dolphins (Lahvis *et al.*, 1995; Reddy *et al.*, 2001; Schwacke *et al.*, 2002; Jaber *et al.*, 2005; Hall *et al.*, 2006). Although organochlorine contamination is decreasing in some areas, levels in Mediterranean Bottlenose Dolphins remain high (Tolosa *et al.*, 1997; Aguilar & Borrell, 2004; Borrell & Aguilar, 2007). Constant monitoring of toxic elements, improvement in developing suitable biomarkers and risk assessment frameworks must be among the priorities for the conservation for such coastal species (Schwacke *et al.*, 2002; Fossi & Marsili 2003; Jaber *et al.*, 2005; Porte *et al.*, 2006).

Various and sometimes high levels of heavy metals have been found in stranded Bottlenose Dolphins from the Mediterranean (e.g. Leonzio *et al.*, 1992; Frodello *et al.* 2000; 2002; Roditi-Elasar *et al.*, 2003; Lahaye *et al.*, 2006). The impact of these metals at the population level is unknown.

3.4 Threat connected especially with migrations

3.5 National and international utilization

4. Protection status and needs

4.1&2 National and International protection status

Cetaceans are protected by law in most Mediterranean countries (Bearzi *et al.*, *in press*). In addition, the existing legislative and treaty instruments in force today provide a potentially solid framework for the conservation of cetaceans in this region.

The western Mediterranean bottlenose dolphin population is listed in Appendix II of the Convention on the Conservation of Migratory Species of Wild Animals (CMS).

The bottlenose dolphin is also listed in Appendix II (Strictly Protected Fauna Species) of the Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention).

This species is fully protected Agreement on the Conservation of Cetaceans in the Black Sea, Mediterranean Sea and contiguous Atlantic area (ACCOBAMS).

Two protocols of the Barcelona Convention address issues of direct relevance to the bottlenose dolphin in the Mediterranean basin. These are: 1) Protocol for the Protection of the Mediterranean Sea against Pollution Resulting from Exploration and Exploitation of the Continental Shelf and the Seabed and its Subsoil (Offshore Protocol): 1994a; 2) Protocol for the Protection of the Mediterranean Sea against Pollution from Land-Based Sources and Activities (LBS Protocol); and 3) Protocol for Specially Protected Areas and Biological Diversity in the Mediterranean (SPA and Biodiversity Protocol).

An other international initiative relevant to Mediterranean bottlenose dolphin protection is the UN Environment Programme (UNEP) Mediterranean Environmental Action Plan - Action Plan for the Conservation of the Cetaceans in the Mediterranean Sea.

In addition, for the Mediterranean European Countries, the following Directives and Regulation are of particular importance in terms of conservation of bottlenose dolphins in the Mediterranean region: 1) Council Directive No 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora (Habitats Directive); 2) Council Regulation (EC) No 1626/94 of 27 June 1994, laying down certain technical measures for the conservation of fishery resources in the Mediterranean Sea; 3) Council Regulation (EC) No 338/97 of 9 December 1996 on the protection of species of wild fauna and flora by regulating trade therein (*Tursiops truncatus* is listed in Appendix II); and 4) Council Regulation (EC) No 812/2004 of 26 April 2004, laying down measures concernine incidental catches of cetaceans in fisheries and amending Regulation (EC) No 88/98.

Tursiops truncatus is listed as “Data Deficient” in the IUCN Red List (Source: <http://www.iucnredlist.org/search/search-basic>); however the Mediterranean bottlenose dolphin, as a whole, has been recently proposed by an IUCN-ACCOBAMS Workshop as “Vulnerable” (Bearzi & Fortuna 2006). This proposal is currently under scrutiny.

4.3 Additional protection needs

Populations of *Tursiops truncatus* in the North and Baltic Seas, western Mediterranean and Black Sea are currently listed in Appendix II of CMS, but not the Bottlenose Dolphins of the eastern Mediterranean Sea. Given the fact that there are not scientific reasons that would suggest to treat the Mediterranean Bottlenose dolphins as divided into two populations, it is believed that this rather peculiar listing was most likely due to a mistake. Alternatively the previous listing could have been affected by the past almost total lack of information on Cetaceans species in the eastern Mediterranean Sea. However, since individuals of *Tursiops truncatus* in the Mediterranean Sea can either be resident, share a wide home range or migrate, it is suggested that all *Tursiops truncatus* populations should be included in App. II of CMS.

4.4 Recent initiatives for increased protection

ACCOBAMS is preparing a Mediterranean Action Plan.

5. Ranges states¹

Listing of states where the occurrence of species has been proved:

ALBANIA, ALGERIA, Bosnia-Herzegovina, CYPRUS, CROATIA, FRANCE, GREECE, ISRAEL, ITALY, Lebanon, LYBIA, MALTA, MONACO, Montenegro, MOROCCO, PORTUGAL, SYRIA, SLOVENIA, SPAIN, TUNISIA, Turkey, UNITED KINGDOM (Gibraltar).

6. Comments from Range States

7. Additional remarks

Preserving populations that are biologically unique and geographically isolated is recognised as an international priority for conservation. The Mediterranean Bottlenose Dolphins are genetically differentiated from those of the Atlantic (Natoli *et al.* 2005). Therefore, the whole Mediterranean bottlenose dolphins should be listed in Appendix II, not only the Western Mediterranean group.

8. References

- Aguilar A., Borrell A. 1994. Abnormally high polychlorinated biphenyl levels in striped dolphins (*Stenella coeruleoalba*) affected by the 1990-1992 Mediterranean epizootic. *Sci. Total Environ.* 154(2-3):237-247.
- Aguilar A., Borrell A. 2004. DDT and PCB reduction in the western Mediterranean from 1987 to 2002, as shown by levels in striped dolphins (*Stenella coeruleoalba*). *Marine Environmental Research* 59:391-404.

¹ CMS Parties in capitals.

- Aguilar A., Borrell A., Reijnders P.J.H. 2002. Geographical and temporal variation in levels of organochlorine contaminants in marine mammals. *Marine Environmental Research* 53:425-452.
- Aguilar A., Raga J.A. 1993. The striped dolphin epizootic in the Mediterranean Sea. *Ambio*, 22(8):524-528.
- Alessandri C., Affronte M., Scaravelli D. 2004. Stomach content analysis of cetaceans stranded along the Italian coasts. *European Research on Cetaceans*, 15:285-288.
- Barone, G. (1895) Modificazioni delle abitudini del delfino comune (*Delphinus delphis*) osservate in Liguria e prodotte dal generalizzarsi della pesca intensiva. *Neptunia*, 10:156-164.
- Barros N.B., Odell D.K. 1990. Food habits of bottlenose dolphins in the southeastern United States. Pages 309-328 in S. Leatherwood and R.R. Reeves, eds. *The bottlenose dolphin*. Academic Press, San Diego.
- Bayed, A. 1997. First data on the distribution of cetacean along the Moroccan coasts. *European Research on Cetaceans* 10:106.
- Bearzi G., Agazzi S., Bonizzoni S., Costa M., Azzellino A. 2007a. Dolphins in a bottle: abundance, residency patterns and conservation of common bottlenose dolphins *Tursiops truncatus* in the semi-closed eutrophic Amvrakikos Gulf, Greece. *Aquatic Conservation: Marine and Freshwater Ecosystems* doi:10.1002/aqc.843.
- Bearzi G., Ferretti S. 2000. Rare report of a bottlenose dolphin foraging in the Venice Lagoon, Italy. *European Research on Cetaceans* 14:110-114.
- Bearzi G., Fortuna C.M. 2006. Common bottlenose dolphin *Tursiops truncatus* (Mediterranean subpopulation). Pp. 64-73 in Reeves R.R., Notarbartolo di Sciara G. (compilers and editors). *The status and distribution of cetaceans in the Black Sea and Mediterranean Sea*. IUCN Centre for Mediterranean Cooperation, Malaga, Spain.
- Bearzi G., Holcer D., Notarbartolo di Sciara G. 2004a. The role of historical dolphin takes and habitat degradation in shaping the present status of northern Adriatic cetaceans. *Aquatic Conservation: Marine and Freshwater Ecosystems* 14:363-379.
- Bearzi G., Notarbartolo di Sciara G., Politi E. 1997. Social ecology of bottlenose dolphins in the Kvarneric (northern Adriatic Sea). *Marine Mammal Science* 13(4):650-668.
- Bearzi G., Politi E., Agazzi S., Azzellino A. 2006. Prey depletion caused by overfishing and the decline of marine megafauna in eastern Ionian Sea coastal waters (central Mediterranean). *Biological Conservation* 127(4):373-382.
- Bearzi G., Politi E., Agazzi S., Bruno S., Costa M., Bonizzoni S. 2005. Occurrence and present status of coastal dolphins (*Delphinus delphis* and *Tursiops truncatus*) in the eastern Ionian Sea. *Aquatic Conservation: Marine and Freshwater Ecosystems* 15:243-257.
- Bearzi G., Politi E., Costa M., Azzellino A., Bastianini M. 2007b. Dolphins in the northern Adriatic Sea: historical data and present occurrence. *Proceedings of Conference "Cetaceans, sea turtles and sharks of the Adriatic Sea"*, 26– 28 October 2006, Cattolica, Italy.
- Bearzi G., Politi E., Notarbartolo di Sciara G. 1999. Diurnal behavior of free-ranging bottlenose dolphins in the Kvarneric (northern Adriatic Sea). *Marine Mammal Science* 15(4):1065-1097.
- Bearzi, G., Fortuna, C.M., Reeves, R.R. In press. Ecology, behaviour and conservation of Common Bottlenose Dolphins *Tursiops truncatus* in the Mediterranean Sea. *Mammal Review*, in press.
- Ben Naceur L., Gannier A., Bradai M.N., Drouot V., Bourreau S., Laran S., Khalfallah N., Mrabet R., Bdioui M. 2004. Recensement du grand dauphin *Tursiops truncatus* dans les eaux tunisiennes. *Bull. Inst. Natn. Scien. Tech. Mer de Salammbô* 31:75-81.

- Birkun A., Kuiken T., Krivokhizhin S., Haines D.M., Osterhaus A.D.M.E., Van de Bildt M.W.G., Joiris C.R., Siebert U. 1998. Epizootic of morbilliviral disease in common dolphins (*Delphinus delphis ponticus*) from the Black Sea. *Vet. Rec.* 144:85-92.
- Blanco C., Salomon O., Raga J.A. 2001. Diet of the bottlenose dolphin (*Tursiops truncatus*) in the western Mediterranean Sea. *Journal of the Marine Biology Association U.K.* 81:1053-1058.
- Blasi, M.F., Pace, D.S. 2006. Interactions between bottlenose dolphin (*Tursiops truncatus*) and the artisanal fishery in Filicudi Island (Italy). Page 114 in I. Kuklik (ed.) Guidelines and abstract book. 20th Annual Conference of the European Cetacean Society, 2-7 April, Gdynia, Poland.
- Bompar, J-M. (2000) Les Cétacés de Méditerranée. Edisud, La Calade, Aix-en-Provence. 186 pp.
- Borrell A., Aguilar A. 2007. Organochlorine concentrations declined during 1987-2002 in western Mediterranean bottlenose dolphins, a coastal top predator. *Chemosphere* 66(2):347-352.
- Borrell A., Aguilar A., Tornero V., Sequeira M., Fernandez G., Alis S. 2006. Organochlorine compounds and stable isotopes indicate bottlenose dolphin subpopulation structure around the Iberian Peninsula. *Environment International* 32(4):516-523.
- Boutiba, Z., Bouderbala, M., Taleb, M.Z. 2003. Bilan actuel de connaissances sur le Cétacé du bassin Algérien, de 1973 à 2003. Pages: 72-76 in "Actes de la 12e Conférence Internationale sur le Cétacés de Méditerranée", RIMMO, Antibes, 15th November 2003.
- Bradai 2001. Le Golfe de Gabes: Ecosystème/Biodiversité. UNEP RAC/SPA: 1-14.
- Briand F., ed. 2000. Fishing down the Mediterranean food webs? CIESM Workshop Series. Kerkyra, Greece. 26-30 July 2000. 99 pp.
- Briand F., ed. 2003. Mediterranean biological time series. CIESM Workshop Monographs 22.
- Briand F., ed. 2004. Investigating the roles of cetaceans in marine ecosystems. CIESM Workshop Monographs 25.
- Brotons, J.M., Munilla, Z., Grau, A.M., Rendell, L.E. 2006. Are pingers effective at reducing interactions between bottlenose dolphins and artisanal fisheries around the Balearic Islands? Page 39 in I. Kuklik (ed.). Guidelines and abstract book. 20th Annual Conference of the European Cetacean Society, 2-7 April, Gdynia, Poland.
- Brunelli, G. 1932. *Biologia industriale dei delfinidi*. Bollettino di Pesca, di Piscicoltura e di Idrobiologia, 3:343-359.
- Cabezón, O., Resendes, A.R., Domingo, M., Raga, J.A., Agusti, C., Alegre, F., Mons, J.L., Dubey, J.P. & Almeria, S. 2004. Seroprevalence of *Toxoplasma gondii* antibodies in wild dolphins from the Spanish Mediterranean coast. *Journal of Parasitology*, 90(3):643-644.
- Caddy J.F. 1997. Mediterranean and Black Sea: FAO Statistical Area 37. FAO Marine Resources Service, Fishery Resources Division. FAO Fisheries Circular 920. FAO, Rome.
- Caddy, J.F., Griffiths, R.C. 1990. Recent trends in the fisheries and environment in the General Fisheries Council for the Mediterranean (GFCM) area. Food and Agriculture Organization, Rome. Studies and Reviews 63:1-71.
- Calzada N., Aguilar A., Sørensen T.B., Lockyer C. 1996. Reproductive biology of female striped dolphin (*Stenella coeruleoalba*) from the western Mediterranean. *J. Zool. Lond.* 240:581-591.
- Cañadas A. 2006. Towards conservation of dolphins in the Alborán Sea. Ph.D. thesis, Universidad Autónoma de Madrid. 351 pp.

- Cañadas A., Hammond P.S. 2006. Model-based abundance estimates for bottlenose dolphins off southern Spain: implications for conservation and management. *Journal of Cetacean Research and Management* 8(1):13-27.
- Cañadas, A., Sagarminaga, R. & García-Tiscar, S. 2002. Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. *Deep Sea Research I*, 49(11):2053-2073.
- Cañadas, A., Sagarminaga, R., Marcos, P., Urquiola, E. 2004. Sector sur (Andalucía y Ceuta). Pag: 133-190 in Raga, J.A., Pantoja, J. (eds.) "Proyecto mediterráneo: zonas de especial interés para la conservación de los cetáceos en el Mediterraneo español". *Naturaleza y Parques Nacionales, Serie técnica*. Ministerio de Medio Ambiente, Organismo autónomo Parques Nacionales.
- Chakroun, F. 1994. Status of cetaceans in Tunisian marine waters. *European Research on Cetaceans* 8:107.
- Commission of the European Communities. 2002. Incidental catches of small cetaceans. Commission Staff Working Paper, Subgroup on Fishery and Environment (SGFEN), Scientific, Technical and Economic Committee for Fisheries (STECF), 83 pages.
- Consiglio, C., Arcangeli, A., Cristo, B., Marini, L., Torchio, A. 1992. Interactions between *Tursiops truncatus* and fisheries along north-eastern coasts of Sardinia, Italy. *European Research on Cetaceans* 6:35-36.
- Convention on the Conservation of European Wildlife and Natural Habitats. 2007. Report of the Standing Committee, 27th meeting, Strasbourg, 26 - 29 November 2007. Document prepared by the Directorate of Culture and Cultural and Natural Heritage, 100 pages.
- Corsolini S., Focardi S., Kannan K., Tanabe S., Borrell A., Tatsukawa R. 1995. Congener profile and toxicity assessment of polychlorinated biphenyls in dolphins, sharks and tuna collected from Italian coastal waters. *Marine Environmental Research* 40(1):33-53.
- Crnkovic, D. 1958. The dolphin problem. *Morsko Ribarstvo*, 10(1):12-14 (in Croatian).
- Cuculic N. 1960. Five-year research programme to advance marine fisheries. *Morsko Ribarstvo* 12(5): 84-86 (in Croatian).
- De Walle, F.B., Nikolopoulou-Tamvakli, M., Heinen, W.J. 1993. Environmental condition of the Mediterranean Sea: European Community Countries. Kluwer Academic Publishers, The Netherlands. 523 pp.
- Deguy, R., Cyrus, J.-L. 1973. Note préliminaire a l'étude des cétacés des côtes Françaises de Méditerranée. *Rev. Trav. Inst. Pêches marit.* 37(2):151-158 (in French).
- Del Rosso R. 1905. *Pesche e pescherie antiche e moderne nell'Etruria marittima*. 2 Volumes. Osvaldo Paggi: Firenze.
- Dhermain F., Ripoll T., Bompar J.M., David L., Di Meglio N. 1999. First evidence of the displacement of a bottle-nosed dolphin *Tursiops truncatus* between Corsica and Hyeres archipelago, south-eastern France. *European Research on Cetaceans* 13:306-311.
- Dhermain, F. 2006. Suivi de la population du Grand Dauphin *Tursiops truncatus* en Corse. Rapport du Groupe d'Etude des Cétacés de Méditerranée pour le compte du Parc National de Port-Cros et du Ministère de l'Ecologie et du Développement durable. 35 pp.
- Di Guardo G., Agrimi U., Morelli L., Cardeti G., Terracciano G., Kennedy S. 1995a. Post mortem investigations on cetaceans found stranded on the coasts of Italy between 1990 and 1993. *Veterinary Records* 136(17):439-442.
- Di Guardo, G., Corradi, A., Agrimi, U., Zizzo, N., Morelli, L., Perillo, L., Kramer, L., Cabassi, E. & Kennedy, S. 1995b. Neuroerinaryathological lesions in cetaceans found stranded from 1991 to 1993 on the coasts of Italy. *European Journal of Veterinary Pathology*, 1:47-51.

- Di Natale A. 1995. Driftnets impact on protected species: observer data from the Italian fleet and proposal for a model to assess the number of cetacean in the by-catch. Col. Vol. Sci. Pap. ICCAT 44(1):255-263.
- Di Natale, A. 1990. Marine mammals interactions in Scombridae fishery activities: the Mediterranean case. Pages 167-174 in the "Report of the GFCM-ICCAT Expert Consultation on Evaluation of Stocks of Large Pelagic Fishes in the Mediterranean Area". FAO Fishery Report No. 449.
- Di Natale, A., Notarbarolo di Sciara, G. 1994. A review of the passive fishing nets and trap fisheries in the Mediterranean Sea and of cetacean bycatch. Rep. Int. Whal. Commn., Special Issue 15:189-202.
- Díaz López B. 2006b. Interactions between Mediterranean bottlenose dolphins (*Tursiops truncatus*) and gillnets off Sardinia, Italy. ICES Journal of Marine Science 63:946-951.
- Díaz López B., Bernal Shirai J.A. 2007. Bottlenose dolphin (*Tursiops truncatus*) presence and incidental capture in a marine fish farm on the north-eastern coast of Sardinia (Italy). J. Mar. Biol. Ass. U.K. 87:113-117.
- Díaz López, B. 2006a. Bottlenose dolphin (*Tursiops truncatus*) predation on a marine fin fish farm: some underwater observations. Aquatic Mammals 32(3):305-310.
- Dubey, J.P., Zarnke, R., Thomas, N.J., Wong, S.K., Van Bonn, W., Briggs, M., Davis, J.W., Ewing, R., Mense, M., Kwok, O.C.H., Romand, S. & Thulliez, P. (2003) *Toxoplasma gondii*, *Neospora caninum*, *Sarcocystis neurona*, and *Sarcocystis canis*-like infections in marine mammals. Veterinary Parasitology, 116:275–296.
- Duguy, R., Casinos, A., Di Natale, A., Filella, S., Ktari-Chakroun, F., Lloze, R., Marchessaux, D. 1983. Répartition et fréquence des mammifères marins en Méditerranée. Rapp. Comm. Int. Mer Médit. 28:223-230.
- Duignan P.J., House C., Odell D.K., Wells R.S., Hansen L.J., Walsh M.T., St.Aubin D.J., Rima B.K., Geraci J.R. 1996. Morbillivirus infection in bottlenose dolphins: evidence for recurrent epizootics in the western Atlantic and Gulf of Mexico. Marine Mammal Science 12(4):499-515.
- Feingold, D., Elasar, M., Goffman, O., Granit, S.-L., Hadar, N., Ratner, E., Scheinin, A., Kerem, D. 2005. Summary of cetacean strandings along the Israeli Mediterranean coast in the past decade (1993-2004). Abstract book of the 19th Annual Conference of the European Cetacean Society, La Rochelle, France, 2-7 April 2005. Pp. 100.
- Fernández-Contreras, M.M., Brotons, J.M., Beltran, C., Aguilar, A. 2002. Interactions between cetaceans and fishing activities in the Balearic Islands. Abstract book of the 16th Annual Conference of the European Cetacean Society, Liege, Belgium, 7-11 April 2002. Pp. 39-40.
- Forcada J., Gazo M., Aguilar A., Gonzalvo J., Fernandez-Contreras M. 2004. Bottlenose dolphin abundance in the NW Mediterranean: addressing heterogeneity in distribution. Mar. Ecol. Prog. Ser. 275:275-87.
- Fortuna C.M. 2006. Ecology and conservation of bottlenose dolphins (*Tursiops truncatus*) in the north-eastern Adriatic Sea. Ph.D. Thesis. University of St. Andrews, UK. 275 pp.
- Fortuna, C.M., Bearzi, G., Notarbartolo di Sciara, G. 1999. Analysis of respiration patterns of bottlenose dolphins observed in the Kvarneric (northern Adriatic Sea, Croatia). European Research on Cetaceans 12:151-155.
- Fortuna, C.M., Wilson, B., Wiemann, A., Riva, L., Gaspari, S., Matesic, M., Oehen, S., Pribanic, S. 2000. How many dolphins are we studying and is our study area big enough? European Research on Cetaceans 14:370-373.
- Fossi M.C., Marsili L. 2003. Effects of endocrine disruptors in aquatic mammals. Pure Appl. Chem. 75(11-12):2235-2247.

- Francesco M., Zucca P., Picciulin M., Zuppa F., Spoto M. 1999. Cetaceans living in the North Adriatic Sea (Gulf of Trieste - Grado Lagoon) - intervention protocol for healthy and distressed animals. *European Research on Cetaceans* 13:410-415.
- Frantzis, A., Alexiasou, P., Paximadis, G., Politi, E., Gannier, A., Corsini-Foka, M. 2003. Current knowledge of the cetacean fauna of the Greek Seas. *J. Cetac. Res. Managt.*, 5: 219-232.
- Frodello J P, Romeo M, Viale D. 2000. Distribution of mercury in the organs and tissues of five toothed-whale species of the Mediterranean. *Environmental Pollution* 108 (3): 447-452
- Frodello J.P., Viale D., Marchand B. 2002. Metal concentrations in the milk and tissues of a nursing *Tursiops truncatus* female. *Marine Pollution Bulletin* 44:551-576.
- Gannier A. 2005. Summer distribution and relative abundance of delphinids in the Mediterranean Sea. *Rev. Écol. (Terre Vie)* 60:223-238.
- Garcia S.M., de Leiva Moreno I., Grainger R. 2005. Global trends in the state of marine fishery resources 1974-2004. Pp. 10-14 in *Review of the State of World Marine Fishery Resources*. FAO Fisheries Technical Paper 457. Food and Agriculture Organization of the United Nations, Rome. 235 pp.
- Gauthier, J.M., Dubeau, H., Rassart, E., Jarman, W.M. & Wells, R.S. 1999. Biomarkers of DNA damage in marine mammals. *Mutation Research*, 444:427-439.
- Gazo M., Fernandez-Contreras M.M., Brotons J.M., Aguilar A. 2004a. Interactions between bottlenose dolphins and artisanal fisheries in the Balearic Islands: May acoustic devices be a solution to the problem? *European Research on Cetaceans* 15:156-159.
- Gazo, M., Forcada, J., Aguilar, A., Fernandez-Contreras, M.M., Borrell, A., Gonzalvo, J., Tornero, V. 2004b. Sector norte (Cataluna e Islas Baleares). Pages 15-65 in Raga, J.A., Pantoja, J. (eds.) "Proyecto mediterraneo: zonas de especial interes para la conservacion de los cetaceos en el Mediterraneo espanol". *Naturaleza y Parques Nacionales, Serie tecnica*. Ministerio de Medio Ambiente, Organismo autonomo Parques Nacionales.
- Genov, T., Kotnjek, P. 2007. Summer distribution of bottlenose dolphins (*Tursiops truncatus*) in Slovenian waters and neighbouring area (northern Adriatic Sea) . 21st Annual Conference of the European Cetacean Society, 22-25 April 2007, San Sebastian, Spain.
- Gnone, G., Nuti, S., Bellingeri, M., Pannoncini, R., Beddocchi, D. 2006. Spatial behaviour of *Tursiops truncatus* along the Ligurian sea coast: preliminary results. *Biol. Mer. Medit.* 13(2):272-273.
- Goffman O., Kerem D., Roditi-Elasar M., Sonin O., Spanier, E. 2001. Bottlenose dolphin-trawler interactions in the Israeli Mediterranean coastline: 1993-2001. IMMRAC / University of Haifa. Unpublished report. 7 pp.
- Gomez de Segura, A., Crespo, E.A., Pedraza, S.N., Hammond, P.S., Raga, J.A. 2006. Abundance of small cetaceans in the waters of the central Spanish Mediterranean. *Marine Biology* 150(1):149-160.
- Gomez de Segura, A., Tomas, J., Raga, J.A. 2004. Sector Centro (Comunidad Valenciana y Region de Murcia). Pages 67-131 in Raga, J.A., Pantoja, J. (eds.) "Proyecto mediterraneo: zonas de especial interes para la conservacion de los cetaceos en el Mediterraneo espanol". *Naturaleza y Parques Nacionales, Serie tecnica*. Ministerio de Medio Ambiente, Organismo autonomo Parques Nacionales.
- Gonzalvo J., Gazo M., Aguilar A. 2004. Distribution and school size patterns of bottlenose dolphins (*Tursiops truncatus*) in northern Spanish Mediterranean waters. *European Research on Cetaceans* 15:386-389.
- Gourret P. 1894. *Les pêcheries et les poissons de la Méditerranée (Provence)*. Librairie J.B. Baillière et Fils: Paris.

- Hadjichristoforou, M. 2004. National implementation report of Cyprus. ACCOBAMS Mop2/Inf 35.
- Hall A.J., McConnell B.J., Rowles T.K., Aguilar A., Borrell A., Schwacke L., Reijnders P.J.H., Wells R.S. 2006. An individual based model framework to assess the population consequences of polychlorinated biphenyl exposure in bottlenose dolphins. *Environmental Health Perspectives* 114(S1):60-64.
- Impetuoso A., Fortuna C.M., Wiemann A., Antolovich W., Holcer D. In press. Presence and distribution of bottlenose dolphins (*Tursiops truncatus*) in the Archipelago of Kornati National Park (Croatia). *European Research on Cetaceans* 17.
- Jaber, J.R., Perez, J., Carballo, M., Arbelo, M., Espinosa de los Monteros, A., Herraiez, P., Munoz, J., Andrada, M., Rodriguez, F. & Fernandez, A. 2005. Hepatosplenic large cell immunoblastic lymphoma in a bottlenose dolphin (*Tursiops truncatus*) with high levels of polychlorinated biphenyl congeners. *Journal of Comparative Pathology*, 132:242–247.
- Kent, R., Leibovitch, M., Goffman, O., Elasar, M., Kerem, D. 2005. Cetacean bycatch in israeli fisheries in the Mediterranean. *ECS* 19:67.
- Lahaye, V., Bustamante, P., Dabin, W., Van Canneyt, O., Dhermain, F., Cesarini, C., Pierce, G.J. & Caurant, F. 2006. New insights from age determination on toxic element accumulation in striped and bottlenose dolphins from Atlantic and Mediterranean waters. *Marine Pollution Bulletin*, 52(10):1219-1230.
- Lahvis G.P., Wells R.S., Kuehl D.W., Stewart J.L., Rhinehart H.L., Via C.S. 1995. Decreased lymphocyte responses in free-ranging bottlenose dolphins (*Tursiops truncatus*) are associated with increased concentrations of PCBs and DDT in peripheral blood. *Environ. Health Persp.* 103(4):62-72.
- Lauriano G., Mackelworth P., Fortuna C.M. 2003. Densità e abbondanza relativa del tursiope (*Tursiops truncatus*) nel parco nazionale dell'Asinara, Sardegna. *Biol. Mar. Medit.* 10(2):717-720.
- Leatherwood, S. 1975. Some observations of feeding behavior of bottle-nosed dolphins (*Tursiops truncatus*) in the northern Gulf of Mexico and (*Tursiops cf. T. gilli*) off southern California, Baja California, and Nayarit, Mexico. *Marine Fisheries Review* 37(9):10-16.
- Leonzio C., Focardi S., Fossi C. 1992. Heavy metals and selenium in stranded dolphins of the Northern Tyrrhenian (NW Mediterranean). *The Science of the Total Environment* 119:77-84.
- Lipscomb T.P., Schulman F.Y., Moffett D., Kennedy S. 1994. Morbilliviral disease in Atlantic bottlenose dolphins (*Tursiops truncatus*) from the 1987-1988 epizootic. *Journal of Wildlife Diseases* 30:567-571.
- Leonart, J. 2005. Mediterranean and Black Sea. Pp. 49-64 in *Review of the State of World Marine Fishery Resources*. FAO Fisheries Technical Paper 457. Food and Agriculture Organization of the United Nations, Rome. 235 pp.
- Manoukian, S., Azzali, M. & Giovagnoli, L. 2004. Features of cetacean groups observed in the Adriatic Sea from 1998 to 2000. *European Research on Cetaceans*, 15:437-441.
- Marelic, D. 1961. Results of the work for the advancement of marine fisheries in 1961. *Morsko Ribarstvo*, 3(1-2):5-7.
- Marini, L., Consiglio, C., Arcangeli, A., Torchio, A., Casale, M., Cristo, B., Nannarelli, S. 1995. Socio-ecology of bottlenose dolphins, *Tursiops truncatus*, along the north-eastern coast of Sardinia (Italy): Preliminary results. *European Research on Cetaceans* 9:139-141.

- Marsili L., Focardi S. 1997. Chlorinated hydrocarbon (HCB, DDTs and PCBs) levels in cetaceans stranded along the Italian coasts: an overview. *Environmental Monitoring and Assessment* 45:129-180.
- Miokovic D., Kovacic D., Pribanic S. 1999. Stomach content analysis of one bottlenose dolphin (*Tursiops truncatus*, Montagu 1821) from the Adriatic Sea. *Natura Croatica* 8(1):61-65.
- Mitra, S., Koutrakis, E., Milani, C. 2004. Cetacean interaction with small scale coastal fisheries: Implications for conservation and damage limitation. *European Research on Cetaceans* 15.
- Mussi B., Miragliuolo A. (2003). I cetacei della costa nord occidentale dell'isola d'Ischia (Canyon di Cuma). Pp. 213-232 in: M. C. Gambi, M. De Lauro and F. Jannuzzi, eds. *Ambiente marino e costiero e territorio delle isole Flegree (Ischia, Procida e Vivara - Golfo di Napoli). Risultati di uno studio multidisciplinare. Memorie dell'Accademia di Scienze Fisiche e Matematiche, Società Italiana di Scienze, Lettere e Arti in Napoli. Liguori Editore. 425 pp.*
- Mussi, B., Gabriele, R., Miragliuolo, A., Battaglia, M. 1998. Cetacean sightings and interactions with fisheries in the archipelago Pontino Campano, southern Tyrrhenian Sea, 1991-1995. *European Research on Cetaceans* 12:63-65.
- Natoli A., Birkun A., Aguilar A., Lopez A., Hoelzel A.R. 2005. Habitat structure and the dispersal of male and female bottlenose dolphins (*Tursiops truncatus*). *Proc. R. Soc. B* (published online, doi:10.1098/rspb.2005.3076).
- Newman, S.J. & Smith, S.A. 2006. Marine mammal neoplasia: a review. *Veterinary Pathology*, 43:865–880.
- Notarbartolo di Sciara G, Venturino MC, Zanardelli M, Bearzi G, Borsani JF, Cavalloni B. 1993. Cetaceans in the central Mediterranean Sea: Distribution and sighting frequencies. *Italian Journal of Zoology* 60:131-138.
- Notarbartolo di Sciara G., Frantzis A., Bearzi G., Reeves R.R. 2006. Sperm whale *Physeter macrocephalus* (Mediterranean subpopulation). Pp. 48-56 in Reeves R.R., Notarbartolo di Sciara G. (compilers and editors). *The status and distribution of cetaceans in the Black Sea and Mediterranean Sea. IUCN Centre for Mediterranean Cooperation, Malaga, Spain.*
- O'Shea T. J., Reeves R.R., Long A.K. 1999. Marine mammals and persistent ocean contaminants. *Proceedings of the Marine Mammal Commission Workshop. Keystone, Colorado, October 12-15, 1998. 150 pp.*
- Orsi Relini L., Cappello M., Poggi R. 1994. The stomach content of some bottlenose dolphins (*Tursiops truncatus*) from the Ligurian Sea. *European Research on Cetaceans* 8:192-195.
- O'Shea T.J., Aguilar A. 2001. Cetaceans and Sirenians. Pp. 427-496 in R.F. Shore and B.A. Rattner (eds). *Ecotoxicology of Wild Mammals. Ecological and Environmental Toxicology Series. John Wiley & Sons Ltd.*
- Öztürk, B., Akan, Y., Topaloglu, B., Keskin, Ç., Karakulak, S., Öztürk, A.A., Dede, A., Türkozan, O. 2004. *Marine life of Turkey in the Aegean and Mediterranean seas. Turkish Marine Research Foundation (TÜDAV) Publications. Marine Education Series. Number 10. 200 pp. Istanbul, Turkey.*
- Öztürk, B., Öztürk, A.A., Dede, A. 2001. Dolphin bycatch in the swordfish driftnet fishery in the Aegean Sea. *Rapp. Comm. int. Mer. Médit.* 36:308.
- Pace, D.S., Pulcini, M., Triossi, F. 1999. *Tursiops truncatus* population at Lampedusa Island (Italy): preliminary results. *European Research on Cetaceans* 15:165-169.
- Peksider-Srica, V. 1931. On the dolphin and its hunting. *Lovacko Ribarski Vjesnik*, 40(10):409-415 (in Croatian).

- Pérez S, De Stephanis R, Verborgh P, Sánchez A. 2006. Absolute abundance of bottlenose dolphins in the Strait of Gibraltar. Poster presented at the 20th Annual Conference of the European Cetacean Society, April 3-6, 2006, Gdynia, Poland.
- Porte, C., Janer, G., Lorusso, L.C., Ortiz-Zarragoitia, M., Cajaraville, M.P., Fossi, M.C. & Canesi, L. 2006. Endocrine disruptors in marine organisms: approaches and perspectives. *Comparative Biochemistry and Physiology Part C*, 143(3):303-315.
- Pulcini M., Triossi F., Pace D.S. 2004. Distribution, habitat use and behaviour of bottlenose dolphins at Lampedusa Island (Italy): results of five years of survey. *European Research on Cetaceans* 15:453-456.
- Pulcini, M., Angradi, A.M., Sanna, A. 1993. Distribution and frequency of cetaceans in the Ligurian-Provencal basin and in the north Tyrrhenian sea (Mediterranean Sea). *European Research on Cetaceans* 7:44-47.
- Raga, J.A., Pantoja, J. (eds.). 2004. Proyecto Mediterraneo: zonas de especial interes para la conservación de los cetáceos en el Mediterraneo español. *Naturaleza y Parques Nacionales, Serie tecnica. Ministerio de Medio Ambiente, Organismo autonomo Parques Nacionales*. 219 pp.
- Raitsos, D.E., Vernicos, D., Northridge, S. 2003. An assessment of damage to fishing nets caused by dolphins in a trammel net fishery in Greece. Abstract book of the 17th Annual Conference of the European Cetacean Society, Las Palmas de Gran Canaria, Spain, 9-13 March 2003. P. 154.
- Reddy, M.L., Reif, J.S., Bachand, A. & Ridgway, S.H. 2001. Opportunities for using Navy marine mammals to explore associations between organochlorine contaminants and unfavorable effects on reproduction. *The Science of the Total Environment*, 274:171-182.
- Reeves, R.R., Smith, B.D., Crespo, E. & Notarbartolo di Sciara, G. (2003) Dolphins, whales, and porpoises: 2000-2010 conservation action plan for the world's cetaceans. IUCN, Gland, Switzerland, 139 pp.
- Reynolds J.E., Lamboeuf M., Ben Abdallah A., Abukhader A., Abdulbari R., Nafati A. 1994. Preliminary findings of the 1993 Libyan marine fisheries frame survey: workshop proceedings. *Technical Assistance in Fisheries Development, Project Libfish. Libfish Technical Briefing Notes 11. FAO, Rome*. 47 pp.
- Ripoll, T., Dhermain, F., Barril, D., Trucchi, R., Bonsignori, B., Roussel, E., David, L., Beaubrun, P. 2004. First summer population estimate of bottlenose dolphins along the north-western coasts of the occidental Mediterranean basin. *European Research on Cetaceans* 15:393-396.
- Roberts C.M., Hawkins J.P. 1999. Extinction risk in the sea. *Trends in Ecology and Evolution* 14(6):241-246.
- Roditi-Elasar M., Karem D., Hornung H., Kress N., Shoham- Frider E., Goffman O., Spanier E. 2003. Heavy metal levels in bottlenose and striped dolphins off the Mediterranean coast of Israel. *Marine Pollution Bulletin* 46:503-512.
- Romeo, T., P. Consoli, G. Florio, M.C. Fossi, C.M. Fortuna, G. Lauriano, F. Andaloro, G. Nortarbartolo di Sciara. 2003. The feluca boat of Messina straits: a platform of opportunity for cetaceans observation. *European Research on Cetaceans* 17.
- Sala E. 2004. The past and present topology and structure of Mediterranean subtidal rocky-shore food webs. *Ecosystems* 7:333-340.
- Salomón, O., Blanco, C. & Raga, J.A. 1997. Diet of the bottlenose dolphin (*Tursiops truncatus*) in the Gulf of Valencia (western Mediterranean). *European Research on Cetaceans*, 11:156.
- Scheinin A., Kerem D., Markovich M., Goffman O., Spanier E. 2005. First summary of coastal sea surveys done off the Mediterranean coast of Israel between the years 1999-

2005. 16th Biennial Conference on the Biology of Marine Mammals, San Diego, California, 12-16 December 2005. Abstract book, p. 249.
- Schulman F.Y., Lipscomb T.P., Moffett D., Krafft A.E., Lichy J.H., Tsai M.M., Taubenberger J.K., Kennedy S. 1997. Histologic, immunohistochemical, and polymerase chain reaction studies of bottlenose dolphins from the 1987-1988 United States Atlantic coast epizootic. *Veterinary Pathology* 34:288-295.
- Schwacke L.H., Voit E.O., Hansen L.J., Wells R.S., Mitchum G.B., Hohn A.A., Fair P.A. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the southeast United States coast. *Environmental Toxicology and Chemistry* 21(12):2752–2764.
- Shane, S. H., R. S. Wells and B. Würsig. 1986. Ecology, behavior and social organisation of the bottlenose dolphin: A review. *Marine Mammal Science* 2:34-63.
- Silvani L., Raich J., Aguilar A. 1992. Bottlenose dolphins, *Tursiops truncatus*, interacting with fisheries in the Balearic Islands, Spain. *European Research on Cetaceans* 6:32–34.
- Stanners, D., Bourdeau, P. 1995. Europe's environment: The Dobris Assessment. European Environment Agency, Copenhagen. 676 pp.
- Stewart, S.E. (2004) Diet analysis of bottlenose dolphin population (*Tursiops truncatus*, Montagu, 1821) in proposed Cres-Losinj Marine Protected Area (MPA), Croatia. M.Sc. thesis, University College of London, U.K.
- Taleb, M.Z., Boutiba, Z. 2003. Régime alimentaire de *Delphinus delphis* et *Tursiops truncatus* du littoral occidental Algérien impact de la pêche sur le comportement alimentaire. Actes de la 12e Conference Internationale sur les cétacés de Méditerranée. Antibes, 15 Novembre 2003. Page 96.
- Tolosa I., Readman J.W., Fowler S.W., Villeneuve J.P., Dachs J., Bayona J.M., Albaiges J. 1997. PCBs in the western Mediterranean. Temporal trends and mass balance assessment. *Deep-Sea Research II* 44(3-4):907–928.
- Tringali, M., Puzzolo, V., and Caltavuturo, G. 2004. A case of opportunistic feeding: the bottlenose dolphin, *Tursiops truncatus*, interference with the European anchovy, *Engraulis encrasicolus*, fishing in the Gulf of Catania (Ionian Sea). *European Research on Cetaceans* 15:105–107.
- Triossi F. and Tizzi R. 2003.? A Cetacean survey in North Adriatic Sea: preliminary results.? Proceedings of the 17th Annual Conference of the European Cetacean Society? (Las Palmas de Gran Canaria, Spain, 9-13 March 2003). In press.
- Trites A.W Christensen V., Pauly D. 1997. Competition between fisheries and marine mammals for prey and primary production in the Pacif Ocean. *Journal of Northwest Atlantic Fishery Science*, 22:173-187.
- Tsur, I., B. Yakobson, D. Elad, D. Moffett, and S. Kennedy, 1997. Morbillivirus infection in a bottlenose dolphin from the Mediterranean. *Sea. Eur. J. Vet. Pathol.* 3, 83–85.
- Tudela S. 2004. Ecosystem effects of fishing in the Mediterranean: an analysis of the major threats of fishing gear and practices to biodiversity and marine habitats. General Fisheries Commission for the Mediterranean, Studies and Reviews 74. 58 pp.
- Tudela, S., Kai Kai, A., Maynou, F., El Andalossi, M. & Guglielmi, P. 2005. Driftnet fishing and biodiversity conservation: the case study of the large-scale Moroccan driftnet fleet operating in the Alboràn Sea (SW Mediterranean). *Biological Conservation* 121:65-78.
- Van Canneyt, O., Peltier, H. (2006) Les echouages de mammiferes marins sur le littoral Français en 2005. CRMM France, September 2006. 45 pp.
- Van de Bildt, M. W., B. E. Martina, B. A. Sidi, and A. D. Osterhaus. 2001: Morbillivirus infection in a bottlenosed dolphin and a Mediterranean monk seal from the Atlantic coast of West Africa. *Vet. Rec.* 148, 210–211.
- Van Bresseem M.-F., I. K. G. Visser, R. L. De Swart, C. Örvell, L. Stanzani, E. Androukaki,

- K. Siakavara and A. D. M. E. Osterhaus. 1993. Dolphin morbillivirus infection in different parts of the Mediterranean Sea. *Archives of Virology* 129(1-4):235-242.
- Vella, A. 1999. Cetacean survey around Maltese islands and sea-user cetacean questionnaire study. *European Research on Cetaceans* 12:66-73.
- Voliani, A., Volpi, C. 1990. Stomach content analysis of a stranded specimen of *Tursiops truncatus*. *Rapp. Comm. int. Mer Médit.* 32:238.
- Wafo, E., Sarrazin, L., Diana, C., Dhermain, F., Schembri, T., Lagadec, V., Pecchia, M. & Rebouillon, P. 2005. Accumulation and distribution of organochlorines (PCBs and DDTs) in various organs of *Stenella coeruleoalba* and a *Tursiops truncatus* from Mediterranean littoral environment (France). *Science of the Total Environment*, 348:115– 127.
- Wilson, B., Reid, R.J., Grellier, K., Thompson, P.M., Hammond. P.S. 2004. Considering the temporal when managing the spatial: a population range expansion impacts protected areas-based management for bottlenose dolphins. *Animal Conservation* 7:331–338.
- Zafiroopoulos, D. & Merlini, L. (2003) A comparative ecological study of bottlenose dolphin (*Tursiops truncatus*) in south and north Evoikos Gulfs. *Proceedings of the 8th International Conference on Environmental Science and Technology Lemnos Island, Greece, 8-10 September 2003.* Vol. B:921-925.
- Zahri, Y., Abid, N., Elouamari, N., Abdellaoui, B. 2004. Étude de l'interaction entre le Grand Dauphin et la pêche à la senne coulissante en Méditerranée Marocaine. *INRH Report*, 52 pages.